

First record of *Hydropsyche spiritoi* Moretti, 1991 in France [Trichoptera, Hydropsychidae]

by Gennaro COPPA¹, Núria BONADA², Thibault DATRY³, Marcos GONZÁLEZ⁴,
Bertrand LAUNAY³, Gwenole LE GUELLEC⁵, Cesc MÚRRIA²
& Carmen ZAMORA-MUÑOZ⁶

¹ 1 rue du Courlis, F - 08350 Villers-sur-Bar, France

² Grup de Recerca Freshwater Ecology and Management (FEM), Departament de Biologia Evolutiva, Ecologia i Ciències Ambientals, Facultat de Biologia, Institut de Recerca de la Biodiversitat (IRBio), Universitat de Barcelona, Diagonal, 643, 08028 Barcelona, Catalonia, Spain

³ Irstea, UR MALY, centre de Lyon-Villeurbanne, 5 rue de la Doua CS70077,
F - 69626 Villeurbanne cedex, France

⁴ Departamento de Zoología, Genética y Antropología Física, Facultad de Biología, Universidad de Santiago de Compostela, Campus Universitario Sur, 15782 Santiago de Compostela, Spain.

⁵ Maison Régionale de l'Eau, boulevard Grisolle, F - 83670 Barjols, France

⁶ Departamento de Zoología, Facultad de Ciencias, Universidad de Granada, Severo Ochoa s/n, 18071, Granada, Spain - E-mail: czamora@ugr.es

Keywords: *Hydropsyche spiritoi*, Trichoptera, new record, Provence-Alpes-Côte d'Azur Region, S-E France.

Several adults and larvae of *H. spiritoi* were collected in the Provence-Alpes-Côte d'Azur Region (S-E France). This material completes our knowledge on the geographical distribution of this species throughout the W-Mediterranean. *H. spiritoi* is a new record for France. Some ecological data and morphological characters are provided.

Première citation d'*Hydropsyche spiritoi* en France [Trichoptera, Hydropsychidae]

Mots clés : *Hydropsyche spiritoi*, Trichoptera, nouvelle citation, Région PACA, S-E France.

Des collectes de Trichoptères dans la Région Provence-Alpes-Côte d'Azur, ont permis de noter pour la première fois la présence d'*H. spiritoi* en France. Ces découvertes complètent nos connaissances sur la distribution géographique de cette espèce dans l'Ouest méditerranéen. Des éléments concernant sa détermination et son écologie sont donnés.

Introduction

Hydropsyche s. l. (Trichoptera, Hydropsychidae) is one of the most diverse Trichopteran genera, with over 600 described species distributed all over biogeographic Regions (MORSE 2016). Larvae of *Hydropsyche* occur in lotic habitats with a wide variety of environmental con-

ditions, frequently reaching high densities (e.g., TANIDA 1980, YAN & LI 2007). In Europe, the genus comprises 120 species (PESI 2016), with 26 recorded in Greece (MALICKY 2005), 25 in France (OPIE-BENTHOS 2016), 23 in Spain (GONZÁLEZ & MARTÍNEZ-MENÉNDEZ 2011, MARTÍNEZ-MENÉNDEZ 2014), 20 in Italy (CIANFICCONI 2002), and 9 in Portugal (GONZÁLEZ & MARTÍNEZ-MENÉNDEZ 2011, MARTÍNEZ-MENÉNDEZ 2014). The Mediterranean countries do not only hold a high number of species but also many endemics. Out of the 74 *Hydropsyche* species recorded in Mediterranean rivers of these countries, 41 are endemic (TIERNO DE FIGUEROA et al. 2013). Moreover, the number of *Hydropsyche* species in these countries is most likely underestimated and new species have been recently described (e.g., GONZÁLEZ & MALICKY 1999, MALICKY 2001, ZAMORA-MUÑOZ et al. 2002, SIPAHILER 2016).

Hydropsyche spiritoi Moretti, 1991 belongs to the *instabilis*-group, originally considered as an endemic element of the Italian Peninsula that was recently cited in Spain (GONZÁLEZ & MARTÍNEZ-MENÉNDEZ 2008). In Italy, it has a wide distribution all over the country (from central Alps to Sicily) but it is more frequently found in the central and southern regions (CIANFICCONI et al. 2006). According to GONZÁLEZ & MARTÍNEZ-MENÉNDEZ (2008), despite the existing adult records in the Aragon region (Spain), *H. spiritoi* is likely more widespread in the north eastern areas. Up to now, *H. spiritoi* has not been recorded in France.

Studied material

Adults and larvae of *H. spiritoi* were collected in the Provence-Alpes-Côte d'Azur Region (S-E France) at the following sites:

Grana River, waterfalls at Saorge, Maritime Alps dept., alt. 430 m: 2 ♂, 17.07.2009 (leg S. Jolivet & G. Coppa).

Cagne River, the confluence of Cagne River and Foussa spring, at Coursegoules, Var dept, alt. 940 m: 1 ♂, 2 ♀, 12.06.2009 (leg M. Derrien & G. Coppa)

Grand Torrent stream, downstream Realtor Lake at La Tour d'Arbois, Aix-en-Provence, Bouches-du-Rhône dept, alt. 129 m: 2 ♂, 24.07.2014.

Bresque River at Sillans-la-Cascade, Var dept., alt. 376 m: 17 ♂, 09.06.2011 and 12.09.2011.

Argens stream at Montaud, near from Chateaufort, Var dept., alt. 223 m: 1 ♂, 08.09.2011.

Vallon-de-Véounes spring at Barjols, Var dept., alt. 240 m: 1 ♂, 26.06.2012.

Vallon de la Combe Amère spring at Salernes, Var dept., alt. 332 m: 1 ♂, 13.10.2011 (leg G. Coppa & G. Le Guellec).

Calavon River, near from Valsaintes, at Simiane-la-Rotonde, Alpes-de-Haute-Provence dept., alt. 500 m: 1 larva, 19.11.2013.

Calavon River, downstream Saut-du-Moine waterfalls, at Simiane-la-Rotonde, Alpes-de-Haute-Provence dept., alt. 467 m: 2 larvae, 19.11.2013, 24.04.2014.

Toulourenc River at Montbrun-les-Bains, Drôme dept., alt. 582 m: 1 larva, 07.01.2015 (leg B. Launay & T. Datry).

Taxonomic remarks

Male adults of *H. spiritoi* are characterized by a combination of the main following morphological characters:

- inferior appendages bearing a very short and thin coxopodite;

- aspect of apical part of the phallic apparatus, which is long and slightly rounded at the tips in lateral view;

- lateroapical projections of the phallic apparatus large and triangular.

Larval gills are lacking on the 7th abdominal segment. Larvae have a submentum with short and wide lateral lobes. The apotome is slightly elongated with parallel-sided lateral margins and with an aboral light spot often joining the light lateral spot forming a V-shape (DE PIETRO 1999). An oral spot can be also present and often joined to the lateral light spot.

Geographical distribution and ecology

Material composed of adults and larvae were collected in the upper basin of karstic temporary streams and rivers. Localities where larvae were collected are characterized by the following environmental data of water: mean temperature, 7.92 ± 6.51 °C; pH, 8.32 ± 0.28 ; conductivity, 575.53 ± 65.54 μ S/cm; dissolved oxygen, concentration (10.78 ± 1.25 mg/l), saturation ($98.02 \pm 1.25\%$). Nevertheless, similar environmental characteristics were reported by GRAF et al. (2008) for *H. spiritoi*.

Based on the present records of *H. spiritoi* from southern France (Provence), the current geographical distribution of this species may cover a larger area throughout the North Western part of the Mediterranean Basin than previously thought. Although biological and ecological traits of *Hydropsyche* larvae are regularly monitored by researchers and water managements (STATZNER & MONDY 2009), more detailed information about its geographical distribution and its morphological variation is still needed to make identifications more reliable. However, additional relevant information on molecular variability of *H. spiritoi* remains necessary and extremely valuable to learn more about its evolutionary origin and its taxonomic position, which could help to assess the status of the existing populations.

Acknowledgements

Thibault Datry & Bertrand Launay were supported by an intermittent river project funded by the "Agence de l'Eau Rhône-Méditerranée-Corse".

References

- CIANFICCONI, F. 2002. The third list of Italian Trichoptera (1990-2000). Pp. 349-358 in *Proceedings of the Xth International Symposium on Trichoptera*. W. Mey (ed).
- CIANFICCONI, F., C. CORALLINI & F. TUCCIARELLI. 2006. Trichoptera endemic to the Italian fauna. Pp. 65-74 in *Proceedings of the XIth International Symposium on Trichoptera*. J. Bueno-Soria et al. (eds).
- DE PIETRO, R. 1999. Identification of the larvae of *Hydropsyche* species from Sicily and peninsular Italy (Trichoptera, Hydropsychidae). *Archiv für Hydrobiologie Supplement* **121** (2): 91-117.
- GONZÁLEZ, M. & H. MALICKY. 1999. Une nouvelle espèce d'*Hydropsyche* du groupe *pellucidula* (Trichoptera, Hydropsychidae). *Braueria*, **26**: 25-26.
- GONZÁLEZ, M. A. & J. MARTÍNEZ-MENÉNDEZ. 2008. Observaciones sobre los tricópteros de la Península Ibérica. X. Tricópteros de Aragón (NE España) (Insecta, Trichoptera). *Boletín Sociedad Entomológica Aragonesa*, **43**: 187-192.
- GONZÁLEZ, M. A. & J. MARTÍNEZ-MENÉNDEZ. 2011. Checklist of the caddisflies of the Iberian Peninsula and Balearic Islands (Trichoptera). *Zoosymposia*, **5**: 115-135.
- GRAF, W., J. MURPHY, J. DAHL, C. ZAMORA-MUÑOZ & M. J. LOPEZ-RODRIGUEZ. 2008. *Distribution and*

- ecological preferences of European freshwater organisms. Volume 1. Trichoptera*. A. Schmidt-Kloiber and D. Hering (Eds). Pensoft, Sofia. Available at: <http://www.pensoft.net/newreleases/14279-toc.html>
- MALICKY, H. 2001. Ein Beitrag zur Kenntnis der Arten der *Hydropsyche instabilis* - Verwandtschaft im östlichen Mittelmeergebiet (Trichoptera, Hydropsychidae). *Linzer Biologische Beiträge*, **33**: 489-518.
- MALICKY, H. 2005. *Die Köcherfliegen Griechenlands (Trichoptera)*. Denisia, 17. Land Oberösterreich Biologiezentrum. Linz.
- MARTÍNEZ-MENÉNDEZ, J. 2014. *Biodiversidad de los tricópteros (Insecta, Trichoptera) de la península ibérica: estudio faunístico y biogeográfico*. PhD thesis, Universidade de Santiago de Compostela, 326 pp.
- MORSE, J. C. (ed.) 2016. *Trichoptera World Checklist*. <http://entweb.clemson.edu/database/trichopt/index.htm> [Accessed 22 June 2016].
- OPIE-BENTHOS (2016). http://www.opie-benthos.fr/opie/pdf/161_pagesdynadocs57062f0c4e1a9.pdf
- PESI (2016). Pan-European Species directories Infrastructure. www.eu-nomen.eu/portal [Accessed 22 June 2016].
- SIPAHLER, F. 2016. Studies on the males of the *Hydropsyche pellucidula* species group in Turkey (Trichoptera, Hydropsychidae). *Nova Acta Científica Compostelana (Biología)*, **23**: 73-83.
- STATZNER, B. & N. MONDY. 2009. Variation of color patterns in larval *Hydropsyche* (Trichoptera): Implications for species identifications and the phylogeny of the genus. *Limnologica*, **39**: 177-183.
- TANIDA, K. 1980. Life history and distribution of three species of *Hydropsyche* (Trichoptera, Hydropsychidae) in the River Kibune (Kyoto, Central Japan), with particular references to the variations in life cycles and the relation of larval growth to their density. *Japanese Journal of Limnology*, **41**: 95-111.
- TIERNO DE FIGUEROA, M., M. J. LÓPEZ-RODRÍGUEZ, S. FENOGLIO, P. SÁNCHEZ-CASTILLO & R. FOCHETTI. 2013. Freshwater biodiversity in the rivers of the Mediterranean Basin. *Hydrobiologia*, **719**: 137-186
- YAN, Y. J. & X. Y. LI. 2007. Abundance, life cycle, production dynamics and trophic basis of a dominant caddisfly (*Hydropsyche carbonaria* McLachlan) in a subtropical stream in China. *Journal of Freshwater Ecology*, **22**: 261-266.
- ZAMORA-MUÑOZ, C., M. A. GONZÁLEZ, J. PICAZO-MUÑOZ & J. ALBA-TERCEDOR. 2002. *Hydropsyche fontinalis*, a new species of the *instabilis*-Group from the Iberian Peninsula (Trichoptera, Hydropsychidae). *Aquatic Insects*, **24**: 189-197.